

Clinical Roles of Interleukin-6 and STAT3 in Oral Squamous Cell Carcinoma

Kenichi Shinagawa¹ · Souichi Yanamoto¹ · Tomofumi Naruse¹ · Akiko Kawakita¹ · Kota Morishita¹ · Yuki Sakamoto¹ · Satoshi Rokutanda¹ · Masahiro Umeda¹

Received: 13 April 2016 / Accepted: 12 October 2016 / Published online: 15 October 2016
© Arányi Lajos Foundation 2016

Abstract The effect inflammation has on cancer prognosis is marked by the presence of cytokines and chemokines. Interleukin-6 (IL-6) is one a multifunctional cytokine that regulates inflammatory responses. We investigated the roles of IL-6 and STAT3 and examined the relationship between IL-6 signaling and clinicopathological factors in patients with oral squamous cell carcinoma (OSCC). We retrospectively examined 116 patients who underwent radical surgery for OSCC. IL-6 and STAT3 expression were detected by immunohistochemistry. IL-6 and STAT3 positivity were detected by IHC, at 78.4 and 80.2 %, respectively. IL-6 expression was significantly associated with pattern of invasion ($P = 0.004$), vascular invasion ($P = 0.003$), and pathological nodal status ($P = 0.019$). Multivariate logistic regression analysis revealed that IL-6 expression was significantly associated with vascular invasion ($P = 0.044$). Meanwhile, there was no significant association between STAT3 expression and clinicopathological factors and no significant relationship between IL-6 and STAT3 expression. IL-6 expression was significantly associated with 5-year disease-free survival. These results suggest that IL-6 is involved in lymphangiogenesis and recurrence in OSCC.

Keywords IL-6 · STAT3 · Oral squamous cell carcinoma · Lymphangiogenesis

Introduction

Oral squamous cell carcinoma (OSCC) accounts for 2–3 % of all cancers and is the 11th most common cancer worldwide [1]. OSCC remains a major cause of morbidity and mortality in patients with head and neck cancer. Despite excellent functional and survival outcomes in patients with early-stage oral cancer, patients with advanced-stage disease continue to have poor survival, with only a 5 % improvement in overall survival in the last 20 years [2]. Therefore, more effective treatment strategies are necessary to improve the survival of patients with OSCC. With advances in tumor immunology, immunotherapy has become an emerging option for cancer therapy [3, 4]. Recently, several studies have reported that biomarkers involved in inflammation and the immune system are useful for understanding the biological behavior of OSCC [5–10].

Previous studies have reported that inflammation affects cancer prognosis [11–19]. IL-6 is a multifunctional cytokine that regulates inflammatory responses [20]. IL-6 plays an important role in many tumor functions, including development, migration, invasion, growth, proliferation, apoptosis, progression, angiogenesis, and differentiation of tumor cells [21]. IL-6 activates the Janus kinase/signal transducer and activator of transcription (JAK/STAT) pathway, phosphatidylinositol-3 kinase (PI3K) pathway, and the mitogen-activated protein kinase (MAPK) pathway [22]. IL-6 binds to IL-6R, which forms a complex with gp130, to activate STAT3 via the JAK/STAT pathway [12, 21, 23]. Activated STAT3 controls proliferation, survival, inflammation, invasion, metastasis, and angiogenesis in normal cells [21, 23, 24]. However, activated STAT3 induces pro-survival and pro-proliferative signaling and contributes to tumor growth of cancer cells [24]. However, few studies have examined the relationship between IL-6 signaling and OSCC biological characteristics. Therefore, we investigated the roles of IL-6 and STAT3 in OSCC and examined

✉ Souichi Yanamoto
syana@nagasaki-u.ac.jp

¹ Department of Clinical Oral Oncology, Unit of Translational Medicine, Graduate School of Biomedical Sciences, Nagasaki University, 1-7-1 Sakamoto, Nagasaki 852-8588, Japan

the relationship between IL-6 signaling and clinicopathological factors of patients with OSCC.

Materials and Methods

Patient Characteristics

We analyzed data from 116 patients with OSCC who had their first visit to our department between April 2008 and March 2013 and were treated, usually with surgery. Ethical approval was obtained from the institutional review board of Nagasaki University Hospital (IRB no.: 15,061,128). Patients who underwent preoperative chemotherapy or radiation therapy or who had insufficient preoperative records were excluded. All patients underwent extensive pretreatment evaluations, including blood chemistry analysis, complete blood count, chest X-ray, computed tomography (CT), and/or magnetic resonance imaging (MRI) of the head and neck area, and thoracoabdominal CT, and staging using the TNM classification of malignant tumors, UICC 7th edition [25].

The pattern of invasion (POI), which was evaluated as types 1–4 as defined previously by Bryne et al. [26], was examined at the host/tumor interface. Perineural invasion was defined as the presence of tumor cells within any of the three layers of the nerve sheath (the epineurium, perineurium, and endoneurium). Vascular invasion was defined as the clear presence of tumor cells within a vascular space (lymphatic space or blood vessel), and the tumor had to be adhered to the vessel endothelium or attached to a thrombus in the vessel.

Immunohistochemistry (IHC)

Biopsy or surgical specimens were taken from the patient with OSCC, formalin-fixed, and paraffin-embedded after surgery. Specimens were cut into 4- μ m serial sections. Serial sections were deparaffinized in xylene, soaked in 10 mM citrate buffer (pH 6), and placed in the autoclave at 121 °C for 5 min for antigenicity activation. Endogenous peroxidase was blocked with 0.3 % H₂O₂ in methanol at room temperature for 30 min. The sections were incubated with primary antibodies (IL6, Abcam, Cambridge, UK; 1:600 dilution and STAT3, 124H6, Cell Signaling Technology, MA, USA; 1:600 dilution) overnight at 4 °C. Immunohistochemical staining was performed with the EnVision system (EnVision+, DAKO, Glostrup, Denmark). Reaction products were visualized using diaminobenzidine (DAB) solution, counterstained with Mayer's hematoxylin, dehydrated, cleared with xylene, and mounted. Immunohistochemistry expression analyses were performed by calculating the sum of distribution scores and intensity scores. The distribution score was defined as the estimated fraction of positively stained tumor area (0: none, 1: <20 %, 2: 20–50 %, 3: 50–80 %, and 4: >80 %). The intensity score

was defined as the estimated staining intensity (0: no staining, 1: weak, 2: moderate, and 3: strong). Total scores ranged from 0 to 7. Score of 5 or more were determined positive expression by Receiver Operating Characteristic (ROC) analysis.

Measurement of C-Reactive Protein (CRP) Level and Neutrophil-to-Lymphocyte Ratio (NLR)

All serum CRP levels and complete blood counts were measured during preoperative examination. NLR values were calculated using neutrophil and lymphocyte counts. The optimal cut-off value for CRP was 3 mg/L, and the NLR cutoff was 2.4, as previously reported [7, 14].

Statistical Analysis

All analyses were performed using SPSS 22.0 (Japan IBM, Tokyo, Japan). Categorical data were assessed by Chi-squared test. Multivariate logistic regression analyses were used to determine the associations between various factors and IL-6 expression. Survival times were calculated from the date of surgery. Disease-free survival (DFS) was defined as the period from the date of surgery to cancer recurrence. Disease-specific survival (DSS) was defined as the period from the date of surgery to death or the end of observation. DFS and DSS of the entire cohort were estimated using the Kaplan-Meier method and compared by log-rank test. Multivariate analyses were assessed using the Cox proportional hazards model. Significant factors from univariate analyses were included in multivariate analysis. *P* values less than 0.05 were considered significant.

Results

Patient Characteristics

Patient demographic characteristics are summarized in Table 1. The mean age at diagnosis was 67.4 years (range, 30–95 years). The male-to-female ratio was 1.27, with 65 male subjects. Diffuse invasion was found in 42 of 116 patients (36.2 %), vascular invasion was found in 54 patients (46.6 %), and perineural invasion was found in 27 patients (23.3 %). Pathological lymph node metastasis was noted in 42 patients (36.2 %). Local recurrence developed in 11 patients (9.5 %) during the follow-up period, which was a mean 48.9 months for all participants (range, 2–88 months).

IL-6 and STAT3 Expression by IHC

IL-6 was strongly expressed in stroma and cancer cell membranes, but it was not expressed in cancer cell cytoplasm (Fig. 1a). STAT3 was strongly expressed in cancer cell

Table 1 Association between IL-6, STAT3, and clinicopathological factors

Characteristic	Number of cases (%)	IL-6			STAT3		
		Negative	Positive	p^a	Negative	Positive	p^a
Age (years)				0.187			0.203
< 65	47(40.5)	13	34		12	35	
≥ 65	69(59.5)	12	57		11	58	
Gender				0.997			0.602
Male	65(56.0)	14	51		14	51	
Female	51(44.0)	11	40		9	42	
Clinical T stage				0.351			0.469
T1–2	92(79.3)	22	70		20	72	
T3–4	24(20.7)	3	21		3	21	
Clinical stage				0.112			0.567
Stage I–II	80(69.0)	21	59		17	63	
Stage III–IV	36(31.0)	4	32		6	30	
POI				0.009			0.520
1–3	74(63.8)	22	52		16	58	
4	42(36.2)	3	39		7	35	
Vascular invasion				0.003			0.741
No	62(53.4)	20	42		13	49	
Yes	54(46.6)	5	49		10	44	
Perineural invasion				0.215			0.307
No	89(76.7)	22	67		20	69	
Yes	27(23.3)	3	24		3	24	
Pathological nodal status				0.032			0.107
No	74(63.8)	21	53		18	56	
Yes	42(36.2)	4	38		5	37	
Local recurrence				0.502			0.294
No	105(90.5)	24	81		19	86	
Yes	11(9.5)	1	10		4	7	
CRP				0.626			0.796
< 3	91(78.4)	21	70		19	72	
≥ 3	25(21.6)	4	21		4	21	
NLR				0.316			0.884
< 2.4	64(55.2)	16	48		13	51	
≥ 2.4	52(44.8)	9	43		10	42	

^a Bolded values indicate $p < 0.05$

POI, Pattern of invasion; CRP, C-reactive protein; NLR, Neutrophil-to-lymphocyte ratio

membranes and nuclei (Fig. 1b). IL-6 and STAT3 positivity rates by IHC were 78.4 and 80.2 %, respectively.

Associations between IL-6 and STAT3 Expression, Clinicopathological Factors, and Survival

IL-6 and STAT3 expression levels are summarized in Table 1. IL-6 expression was significantly associated with POI, vascular invasion, and pathological nodal status. Multivariate logistic regression analysis revealed that IL-6 expression was significantly associated with vascular invasion ($P = 0.044$)

(Table 2). Meanwhile, there was no significant association between STAT3 expression and clinicopathological factors. There was no significant relationship between IL-6 and STAT3 expression (Table 3). Univariate analyses by log-rank test revealed that POI ($P < 0.001$), vascular invasion ($P < 0.001$), perineural invasion ($P < 0.001$), pathological nodal status ($P < 0.001$), and IL-6 (Fig. 2, $P = 0.010$) were significantly associated with 5-year DFS (Table 4). Predictors associated with 5-year DFS in univariate analyses were included in the Cox proportional hazards model, and this multivariate analysis showed that POI and pathological nodal status




Fig. 1 Representative immunohistochemical staining for IL-6 and STAT3 in OSCC. **a** IL-6 was strongly positive in cell stroma and on cancer cell membranes (intensity score 3), but there was no IL-6 staining in the cancer cell cytoplasm. **b** STAT3 was strongly expressed in cancer cell membranes and nuclei (intensity score 3)

were independent predictors of 5-year DFS. Univariate analyses by log-rank test revealed that clinical stage ($P = 0.032$), POI ($P < 0.001$), vascular invasion ($P = 0.026$), perineural invasion ($P = 0.004$), and pathological nodal status ($P < 0.001$) were significantly associated with 5-year DSS. According to the Cox proportional hazards multivariate analysis, pathological

Table 2 Association between IL-6 expression and clinicopathological factors by multivariate analysis

Characteristics	IL-6		
	HR	CI	P
POI			0.114
1–3	Reference		
4	3.84	0.73–20.3	
Vascular invasion			0.044
No	Reference		
Yes	3.30	1.03–10.6	
Pathological nodal status			0.945
No	Reference		
Yes	0.95	0.19–4.70	

Bolded values indicate $p < 0.05$

POI, Pattern of invasion; HR, hazard ratio; CI, confidence interval

Table 3 Relationship between IL-6 expression and STAT3 expression

	STAT3 expression		$P = 1.000$
	Negative (n = 23)	Positive (n = 93)	
IL-6 expression			
Negative (n = 25)	5	20	
Positive (n = 91)	18	73	

nodal status was the only independent predictor of 5-year DSS.


Discussion

IL-6 is a multifunctional cytokine that regulates immune responses [20]. IL-6 production is increased in response to various stimuli, such as infection and inflammation. Cancer stimulates inflammation that can lead to increased expression of IL-6-modulating factors, such as IL-1 β , COX-2, PGE2, and TGF- β [22]. The IL-6 receptor is a type I cytokine receptor complex that consists of the ligand-binding IL-6Ra chain (also called CD126) and the gp130 (also called CD130), which is responsible for downstream signaling [21]. IL-6 activates the JAK/STAT, PI3K, and the MAPK pathways [8, 22]. This signaling influences cell migration, malignant tumor growth and invasion, anti-apoptotic signaling, angiogenesis, and bone remodeling [8–10, 22, 27, 28].

Shinriki et al. demonstrated that IL-6 signaling via the PI3K-Akt pathway stimulates VEGF-C synthesis and lymphangiogenesis in OSCC [27]. VEGF-C is an essential chemotactic and survival factor during embryonic and inflammatory lymphangiogenesis [29]. Furthermore, several studies have reported that the VEGF-C is associated with lymph node metastasis, and its expression is a prognostic factor for various cancers including oral cancer [30–32]. In this study, we examined the association between IL-6 expression and clinicopathological factors in patients with OSCC by IHC. Our data indicate that IL-6 expression is associated with POI, vascular invasion, and pathological nodal status in OSCC. In particular, vascular invasion strongly correlated with IL-6 expression, suggesting that IL-6 is involved in lymphangiogenesis in OSCC.

On the other hand, several studies have reported that IL-6 activates the JAK/STAT3 pathway in various cancers [12, 15, 24, 28]. Yadav et al. demonstrated that IL-6 promotes epithelial-mesenchymal transition (EMT) changes via the JAK/STAT3 pathway [28]. STAT3 is downstream mediator of IL-6 and represses E-cadherin transcription via the snail-related zinc-finger transcription factor [28]. E-cadherin loss in tumor cells is a widely known hallmark of EMT. Additionally, some

Fig. 2 Kaplan-Meier curves for 5-year disease-free survival according to IL-6 expression in patients with OSCC. The IL-6 overexpression group had a significantly lower disease-free survival rate than that of the low expression group ($P = 0.010$)



studies have reported that IL-6 overexpression plays transcriptional and regulatory roles in invasion and metastasis, leading to poor prognoses for patients with head and neck cancers [9, 17, 22, 24, 28]. In this study, however, the relationship between IL-6 and STAT3 expression was not significant. In addition, STAT3 was not significantly associated with clinicopathological factors. Therefore, our results suggested that IL-6 promotes lymphangiogenesis via the PI3K-Akt pathway rather than the JAK/STAT3 pathway.

Some studies reported that post-operative serum or salivary IL-6 levels could predict recurrence and survival in patients with head and neck cancer [9, 10, 33]. Sato et al. indicated that post-operative salivary IL-6 level is a useful marker for OSCC locoregional recurrence [10]. In this study, IL-6 expression was significantly associated with 5-year DFS, suggesting that IL-6 evaluation by IHC could predict time to recurrence, including locoregional recurrence and distant metastasis. However, any correlation between IL-6 expression and local recurrence was not clear, leading us to conclude that IL-6 was primarily involved in lymphangiogenesis and metastasis of OSCC.

CRP and NLR are indicators of inflammatory response and often checked at during preoperative examination. Tumor growth can cause tissue inflammation and an immune response to tumor antigens [11, 12]. Furthermore, IL-6 secreted by cancer cells induces CRP production, which is a tumor antigen that links chronic inflammation and tumor growth [14–17]. Leukocyte count is usually increased in response to infection, inflammation, allergic reaction, and malignancy. Although neutrophils have anti-tumor effects, lymphocytes are most responsible for controlling cancer progression, and increasing NLR is less efficacious [18].

Therefore, NLR elevation has been suggested to be associated with poor prognosis. Some studies report a relationship between prognosis and both high preoperative CRP levels and NLR [5, 7, 19]. Tumor cells produce various cytokines and chemokines that induce inflammatory responses, including a diverse leukocyte population, cytokines, and cytotoxic mediators [11]. This study found no relationship between IL-6 or STAT3 expression and increasing preoperative CRP levels or NLR in OSCC. In our study, the proportion of patients with high CRP levels was small compared with that of other reports [5, 7]. In cases of oral cancer, when the differential diagnosis includes acute dental infection, antibiotics are administered early. Inflammation-related oral cancer becomes suspected when chronic inflammation remains, with no CRP level elevation. Some studies reported that elevated NLR is associated with worse prognosis [7, 19]. Neutrophils induced by GCSF from the tumor aid in tumor progression by producing chemical mediators, such as VEGF, and causing a relative reduction in lymphocyte number and a decrease in lymphocyte-dependent cellular immune reactions [18, 34, 35]. Because lymphocyte-mediated cellular immune responses are reduced when NLR is high, high NLR is often seen in tumor progression and is a predictor of poor prognosis [18]. In this study, the relationship between IL-6 and STAT3 expression and increased preoperative NLR was not significant. Recent studies have reported that elevated NLR could be used to predict response to neoadjuvant chemoradiation and prognosis in rectal cancer [36, 37]. Therefore, if patients with lower baseline NLR values underwent neoadjuvant chemoradiation, they would be more likely to have better responses. We could not evaluate the relationship between NLR value and neoadjuvant therapy efficacy

Table 4 Univariate and multivariate analyses of risk factors influencing survival

Characteristics	Disease-free survival				Disease-specific survival			
	Univariate		Multivariate		Univariate		Multivariate	
	5-year DFS	<i>p</i> ^a	HR (95 % CI)	<i>p</i> ^a	5-year DSS	<i>p</i> ^a	HR (95 % CI)	<i>p</i> ^a
Age (years)		0.424	-	-		0.332	-	-
< 65	65.7 %				75.4 %			
≥ 65	62.2 %				87.0 %			
Gender		0.203	-	-		0.329	-	-
Male	70.6 %				83.5 %			
Female	55.5 %				81.2 %			
Clinical T stage		0.065	-	-		0.588	-	-
T1–2	67.7 %				83.5 %			
T3–4	50.0 %				76.6 %			
Clinical stage		0.050	-	-		0.032		0.753
Stage I–II	71.6 %				88.6 %		Reference	
Stage III–IV	46.6 %				61.7 %		1.20 (0.38–3.80)	
POI		<0.001		0.002		<0.001		0.227
1–3	90.1 %		Reference		94.6 %		Reference	
4	17.2 %		5.90 (1.97–17.7)		60.7 %		3.03 (0.50–18.3)	
Vascular invasion		<0.001		0.642		0.026		0.746
No	78.0 %		Reference		89.7 %		Reference	
Yes	47.7 %		1.19 (0.58–2.43)		73.7 %		1.22 (0.36–4.14)	
Perineural invasion		<0.001		0.612		0.004		0.572
No	75.2 %		Reference		87.3 %		Reference	
Yes	26.6 %		0.84 (0.42–1.67)		65.9 %		0.71 (0.22–2.33)	
Pathological nodal status		<0.001		0.004		<0.001		0.031
No	91.5 %		Reference		96.8 %		Reference	
Yes	17.2 %		5.11 (1.70–15.4)		56.0 %		14.0(1.27–154)	
CRP		0.522	-	-		0.066	-	-
< 3	62.0 %				87.5 %			
≥ 3	70.7 %				65.4 %			
NLR		0.482	-	-		0.351	-	-
< 2.4	65.8 %				78.1 %			
≥ 2.4	62.0 %				86.6 %			
IL-6		0.010		0.268		0.126	-	-
Negative	88.0 %		Ref.		95.8 %			
Positive	56.5 %		2.29 (0.53–9.95)		77.9 %			
STAT3		0.338	-	-		0.184	-	-
Negative	72.8 %				93.8 %			
Positive	61.7 %				78.7 %			

^a Bolded values indicate *p* < 0.05

CI confidence interval, POI Pattern of invasion, CRP C-reactive protein, NLR Neutrophil-to-lymphocyte ratio

because patients who underwent neoadjuvant chemoradiation or chemotherapy were not included in our cohort.

In conclusion, our data indicate that IL-6 expression is associated with POI, vascular invasion, and pathological nodal status in OSCC. In particular, vascular invasion strongly correlated with IL-6 expression but not STAT3. Therefore,

the present study suggests that IL-6 involved in lymphangiogenesis and recurrence in OSCC. Further studies to elucidate the mechanism of IL-6-mediated VEGF-C synthesis may provide information leading to the development of treatment targeting the PI3K-Akt pathway and improved outcomes in patients with OSCC.

Acknowledgments This study was supported in part by Grant 15 K11295 from the Ministry of Education, Culture, Sports, Science and Technology in Japan to S. Yanamoto. We would like to thank Editage (<http://www.editage.jp/>) for English-language editing.

Compliance with Ethical Standards

Conflict of Interest The authors have no conflicts of interests to declare in this study.

References

- Fitzmaurice C, Dicker D, Pain A et al (2015) The global burden of cancer 2013. *JAMA Oncol* 1:505–527
- Rogers SN, Brown JS, Woolgar JA et al (2009) Survival following primary surgery for oral cancer. *Oral Oncol* 45:201–211
- Turcotte S, Rosenberg SA (2011) Immunotherapy for metastatic solid cancers. *Adv Surg* 45:341–360
- Bell RB, Leidner R, Feng Z et al (2015) Developing an immunotherapy strategy for the effective treatment of oral, head and neck squamous cell carcinoma. *J Oral Maxillofac Surg* 73:S107–S115
- Chen HH, Chen IH, Liao CT et al (2011) Preoperative circulating C-reactive protein levels predict pathological aggressiveness in oral squamous cell carcinoma: a retrospective clinical study. *Clin Otolaryngol* 36:147–153
- Huang SF, Wei FC, Liao CT et al (2012) Risk stratification in oral cavity squamous cell carcinoma by preoperative CRP and SCC antigen levels. *Ann Surg Oncol* 19:3856–3864
- Fang HY, Huang XY, Chien HT et al (2013) Refining the role of preoperative C-reactive protein by neutrophil/lymphocyte ratio in oral cavity squamous cell carcinoma. *Laryngoscope* 123:2690–2699
- Culig Z (2013) Interleukin-6 as a therapy target in oral squamous carcinoma. *Expert Opin Ther Targets* 17:53–59
- Jinno T, Kawano S, Maruse Y et al (2015) Increased expression of interleukin-6 predicts poor response to chemoradiotherapy and unfavorable prognosis in oral squamous cell carcinoma. *Oncol Rep* 33:2161–2168
- Sato J, Ohuchi M, Wada M et al (2015) Differences in sequential posttreatment salivary IL-6 levels between patients with and patients without locoregional recurrences of oral squamous cell carcinoma: part III of a cohort study. *Oral Surg Oral Med Oral Pathol Oral Radiol* 120:751–760
- Coussens LM, Werb Z (2002) Inflammation and cancer. *Nature* 420:860–867
- Yu H, Kortylewski M, Pardoll D (2007) Crosstalk between cancer and immune cells: role of STAT3 in the tumour microenvironment. *Nat Rev Immunol* 7:41–51
- Heikkilä K, Ebrahim S, Lawlor DA (2007) A systematic review of the association between circulating concentrations of C reactive protein and cancer. *J Epidemiol Community Health* 61:824–833
- Allin KH, Bojesen SE, Nordestgaard BG (2009) Baseline C-reactive protein is associated with incident cancer and survival in patients with cancer. *J Clin Oncol* 27:2217–2224
- Yu H, Pardoll D, Jove R (2009) STATs in cancer inflammation and immunity: a leading role for STAT3. *Nat Rev Cancer* 9:798–809
- Kruse AL, Luebbbers HT, Gratz K et al (2010) C-reactive protein levels: a prognostic marker for patients with head and neck cancer? *Head Neck Oncol* 2:21
- Neurath MF, Finotto S (2011) IL-6 signaling in autoimmunity, chronic inflammation and inflammation-associated cancer. *Cytokine Growth Factor Rev* 22:83–89
- Paramanathan A, Saxena A, Morris DL (2014) A systematic review and meta-analysis on the impact of pre-operative neutrophil lymphocyte ratio on long term outcomes after curative intent resection of solid tumours. *Surg Oncol* 23:31–39
- Perisanidis C, Kornek G, Poschl PW et al (2013) High neutrophil-to-lymphocyte ratio is an independent marker of poor disease-specific survival in patients with oral cancer. *Med Oncol* 30:334
- de Oliveira MV, Fraga CA, Gomez RS et al (2009) Immunohistochemical expression of interleukin-4, -6, -8, and -12 in inflammatory cells in surrounding invasive front of oral squamous cell carcinoma. *Head Neck* 31:1439–1446
- Guo Y, Xu F, Lu T et al (2012) Interleukin-6 signaling pathway in targeted therapy for cancer. *Cancer Treat Rev* 38:904–910
- Tawara K, Oxford JT, Jorcyk CL et al (2011) Clinical significance of interleukin (IL)-6 in cancer metastasis to bone: potential of anti-IL-6 therapies. *Cancer Manag Res* 3:177–189
- Yoshimura A (2006) Signal transduction of inflammatory cytokines and tumor development. *Cancer Sci* 97:439–447
- Mali SB (2015) Review of STAT3 (signal transducers and activators of transcription) in head and neck cancer. *Oral Oncol* 51:565–569
- Sobin LH, Gospodarowicz MK, Wittekind C (2009) International Union against Cancer (UICC) TNM classification of malignant tumors. 7th ed. Wiley-Blackwell, Oxford
- Bryne M, Koppang HS, Lilleng R et al (1992) Malignancy grading of the deep invasive margins of oral squamous cell carcinomas has high prognostic value. *J Pathol* 166:375–381
- Shinriki S, Jono H, Ueda M et al (2011) Interleukin-6 signaling regulates vascular endothelial growth factor-C synthesis and lymphangiogenesis in human oral squamous cell carcinoma. *J Pathol* 225:142–150
- Yadav A, Kumar B, Datta J et al (2011) IL-6 promotes head and neck tumor metastasis by inducing epithelial-mesenchymal transition via the LAK-STAT3-SNAIL signaling pathway. *Mol Cancer Res* 9:1658–1667
- Otrock ZK, Makarem JA, Shamseddine AI (2007) Vascular endothelial growth factor family of ligands and receptors: review. *Blood Cells Mol Dis* 38:258–268
- Omoto I, Matsumoto M, Okumura H et al (2014) Expression of vascular endothelial growth factor-C and vascular endothelial growth factor receptor-3 in esophageal squamous cell carcinoma. *Oncol Lett* 7:1027–1032
- Arinaga M, Noguchi T, Takeno S et al (2003) Clinical significance of vascular endothelial growth factor C and vascular endothelial growth factor receptor 3 in patients with nonsmall cell lung carcinoma. *Cancer* 97:457–464
- Wakisaka N, Hirota K, Kondo S et al (2012) Induction of lymphangiogenesis through vascular endothelial growth factor-C/vascular endothelial growth factor receptor 3 axis and its correlation with lymph node metastasis in nasopharyngeal carcinoma. *Oral Oncol* 48:703–708
- Duffy SA, Taylor JMG, Terrell JE et al (2008) Interleukin-6 predicts recurrence and survival among head and neck cancer patients. *Cancer* 113:750–757
- Jablonska J, Leschner S, Westphal K et al (2010) Neutrophils responsive to endogenous IFN-beta regulate tumor angiogenesis and growth in a mouse tumor model. *J Clin Invest* 120:1151–1164
- Li MX, Liu XM, Zhang XF et al (2014) Prognostic role of neutrophil-to-lymphocyte ratio in colorectal cancer: a systematic review and meta-analysis. *Int J Cancer* 134:2403–2413
- Shen L, Zhang H, Liang L et al (2014) Baseline neutrophil-lymphocyte ratio (≥ 2.8) as a prognostic factor for patients with locally advanced rectal cancer undergoing neoadjuvant chemoradiation. *Radiat Oncol* 9:295
- Kim IY, You SH, Kim YW (2014) Neutrophil-lymphocyte ratio predicts pathologic tumor response and survival after preoperative chemoradiation for rectal cancer. *BMC Surg* 14:94